An incision is the cut in your skin made during your surgery.

The incision may be sewn closed (stitches), glued, stapled, or taped.

Taking care of your incision after your surgery is important to reducing the chance of getting an infection and helping you to return to normal sooner.



To listen to the instructions: Scan this QR Code

Dressing Change/ Incision Care



Instructions and Q&A



Can I take a bath when I get home?

- DO NOT take a bath, sit in a hot tub, or get into a swimming pool.
- DO NOT soak your incision in water.
- YOU CAN take a shower beginning two days after your surgery.

When should I remove the dressing?

- REMOVE the dressing two (2) days after your surgery. Steps 3 and 4 are instructions for removing the dressing, cleaning your incision and replacing your dressing.
- DO NOT REMOVE any small strips across your incision ("Steri Strips"). Let them fall off on their own.

What are the steps I should follow to remove the dressing?

- CLEAN YOUR HANDS either by washing your hands with soap and water or using waterless hand sanitizer.
- LOOSEN THE TAPE around the dressing using alcohol on a cotton ball to dab the edges of the tape until it comes loose.
- NEVER tug on or rip off tape or dressing.
- REMOVE THE DRESSING slowly. If the dressing sticks to your incision put clean warm water on a cotton ball and dab the dressing until it comes loose. Never pull on the dressing.
- CHECK the dressing. If the dressing has a foul smell or if pus is coming from your incision, call your doctor.
- DISCARD the old dressing in the trash and clean your hands again.

My doctor told me to clean my incision and then to replace the dressing. How do I do that?

 TO CLEAN YOUR INCISION and REPLACE YOUR DRESSING gather a

- clean, wash rag/cloth a bath towel, and a clean dressing/gauze and tape. The wash rag/cloth and bath towel should be freshly washed.
- CLEAN YOUR INCISION EVERY OTHER DAY using plain soap on the clean, wet wash rag/cloth. Gently soap your incision to remove any old blood or drainage. Do not scrub your incision.
- RINSE YOUR INCISION using the rinsed wash rag/cloth, gently remove the soap.
 Do not rub hard.
- PAT YOUR INCISION DRY using the clean bath towel.
- DO NOT USE alcohol, peroxide, or iodine to clean your incision.
- DO NOT APPLY lotions, creams, or herbal remedies to your incision.
- PLACE a clean dressing/gauze over your incision, being careful not to touch your incision. Secure the dressing/gauze with tape.

How do I manage my incision when taking a shower?

- BEFORE EACH SHOWER gather a clean wash rag/cloth, a bath towel, underclothes (underwear and bra) and clothes. These should be freshly washed.
- TAKE A SHOWER, not a bath, beginning two days after your surgery.
- KEEP YOUR DRESSING DRY by covering the dressing and the area around the dressing with a clean plastic bag or clear, plastic wrap.
- REMOVE the plastic bag or clear, plastic wrap after your shower and discard.
- CHANGE YOUR DRESSING AGAIN if you notice that the dressing has gotten damp or wet during your shower.
- WEAR newly washed underclothes (underwear, bra) and clothes.

6 What if I cannot shower?

- TAKE A SPONGE BATH every day beginning two days after you get home.
- BEFORE EACH SPONGE BATH gather a clean wash rag/cloth, a bath towel, underclothes (underwear and bra) and clothes. These should be freshly washed.
- FOR YOUR SPONGE BATH sit in a chair.
 Use plain soap on a clean, wet wash rag/
 cloth to gently soap your body starting
 with your face, then your neck, chest and
 back, arms, underarms, legs and then
 your private parts. Rinse the wash rag/
 cloth and then wipe the soap off of your
 body.
- DRY YOUR BODY using the clean bath towel.
- WEAR newly washed underclothes (underwear, bra) and clothes.

What should I do if my incision bleeds?

- PLACE a clean, dry dressing/gauze over the area that is bleeding.
- APPLY gentle pressure for a two (2) to three (3) minutes. Do not push too hard.
- The bleeding should stop after two (2) to three (3) minutes.

Mhen to call the doctor?

- If the bleeding does not stop after you follow the instructions listed on number 7 above.
- If your incision breaks open.
- If the redness around your incision increases or spreads more than a half of inch from your incision.
- If you see pus coming from your incision.
- If you get a temperature of 100.5 degrees or higher.